Indicators	Srikakulam				Vizianagaram				Visakhapatnam			
	DLHS-4 (2012-13)		DLHS-3 (2007-08)		DLHS-4 (2012-13)		DLHS-3 (2007-08)		DLHS-4 (2012-13)		DLHS-3 (2007-08)	
	TOTAL	RURAL	TOTAL	RURAL	TOTAL	RURAL	TOTAL	RURAL	TOTAL	RURAL	TOTAL	RURAL
Sample Size												
Ever married women age15-49 years	1023	518	861	744	949	458	1047	865	1356	748	1088	676
Population and household profile												
Sex ratio at birth (Male per 100 Female)	90.97	87.0	94.0	94.7	94.0	93.1	95.9	96.0	98.1	98.2	99.2	100.0
Percentage of households												
Improved source of drinking water[1]	84.7	73.7	70.4	68.6	97.5	95.2	84.1	81.3	95.7	93.3	73.9	57.8
Having access to improved toilet facility	54.5	23.7	18.5	12.4	50.7	25.5	18.5	8.0	59.8	29.5	42.6	13.7
Marriage												
Mean age at marriage for girls (marriages that occurred during the reference period)	21.1	20.0	18.5	18.4	20.07	19.53	18.8	18.4	20.1	19.2	19.7	18.7
Mean age at marriage for boys (marriages that occurred during the reference period)	26.3	25.6	24.4	24.4	25.14	24.95	24.3	23.3	25.0	24.2	23.7	21.9
Percentage of currently married women married below age 18 years (marriages that occurred during the reference period)	4.3	7.5	31.5	32.8	11.4	14.8	28.6	30.8	17.8	21.0	22.6	28.7
Characteristics of women (%)												
Currently married women who are illiterate	37.7	53.9	55.8	59.2	41.4	53.6	57.4	63.1	45.5	61.3	49.4	69.0
Currently married women with 10 or more years of schooling	38.0	24.2	10.5	7.4	31.4	18.2	10.9	6.6	34.2	19.4	21.6	9.0
Fertility (%)												
Births to women aged 15-19 years out of total births[3]	3.8	5.6	18.4	18.2	7.3	3.9	21.5	22.8	3.2	5.0	15.9	17.9
Women aged 20-24 years reporting birth order of 2 & above	47.3	53.5	56.5	56.6	43.1	39.5	55.0	57.1	56.9	61.9	56.5	58.1
Women aged 15-49 years who reported birth order of 3 & above	10.3	10.4	12.0	10.8	2.1	2.4	16.4	19.1	15.0	22.2	20.1	25.6
Women with two children wanting no more children	58.3	35.3	21.8	21.2	35.0	50.0	21.4	20.1	35.3	30.8	24.2	21.7
Mean no. of children ever born to women age 40-49 years	2.3	2.3	3.2	3.2	2.5	2.6	3.3	3.4	2.5	2.7	3.1	3.5
Current use of Family Planning Methods (%)												
Any method	66.9	68.8	74.2	74.1	79.1	79.9	70.4	70.8	79.1	79.9	66.6	60.3
Any modern method	66.7	68.8	73.8	74.1	78.9	79.9	70.0	70.5	73.3	69.6	66.5	60.3
Female sterilization	63.4	67.0	70.3	71.3	75.8	78.0	62.5	64.1	66.9	62.0	45.4	36.9
Male sterilization	2.8	2.2	2.7	2.2	2.1	1.2	6.1	5.6	5.3	6.5	18.6	23.0
Pill	0.4	0.2	0.1	0.0	0.1	0.2	0.4	0.4	0.2	0.1	0.1	0.0
IUD	0.5	0.2	0.1	0.1	0.7	0.2	0.4	0.3	NA	NA	0.7	0.0
Condom	0.2	NA	0.6	0.4	0.2	NA	0.6	0.3	1.3	1.3	1.6	0.3
Any Traditional method	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Unmet Need for Family Planning (%)												
Total unmet need[4]	21.3	18.6	5.9	5.5	11.2	11.1	6.9	6.7	13.7	14.3	10.2	11.7
Unmet Need for spacing	9.7	10.0	3.3	3.3	10.6	11.1	4.4	4.3	10.7	11.8	5.4	6.5
Unmet Need for limiting	11.6	8.6	2.6	2.2	0.6	0.0	2.5	2.4	3.0	2.6	4.8	5.2
Quality of Family Planning Services (%)												
Non-users ever advised by health personnel to adopt any family planning method	37.1	36.5	14.5	14.6	21.2	12.2	14.5	14.6	23.1	19.4	13.0	13.8
Current users told about side effects of method	4.3	5.3	6.7	10.0	10.5	10.7	11.1	10.0	12.3	14.6	26.9	50.0
Users who received follow-up services for sterilization and IUD within 48 hours	91.7	95.6	97.7	97.8	58.0	53.6	97.7	97.8	84.1	85.9	90.4	85.2
Post-partum adoption of Family Planning for sterilization	97.9	98.7	89.5	90.7	97.9	98.7	86.1	86.9	96.3	97.4	79.6	76.3
Antenatal Care[5] (Women who had last live/still birth during reference period) (%)												
Pregnant women who received any antenatal check-up	98.3	98.9	98.6	98.9	98.9	99.0	90.8	89.1	85.7	77.1	89.2	84.8
Pregnant women who had antenatal check-up in first trimester	83.4	80.6	66.6	62.6	91.9	93.8	57.8	52.1	86.5	86.5	59.7	51.4
Pregnant women who had three or more ANC visits	92.6	92.5	92.6	94.0	96.0	94.8	82.6	82.3	96.2	93.8	76.4	74.0
Pregnant women who had at least one tetanus toxoid injection	94.3	97.8	92.9	92.5	98.3	99.0	86.5	83.6	98.1	98.0	85.4	79.4
Pregnant women whose Blood Pressure (BP) is taken	94.9	93.6	98.4	98.7	88.4	89.7	89.8	87.7	95.6	91.9	88.1	82.5
Pregnant women who had blood tested (Hb)	89.8	90.3	98.4*	98.8*	88.4	89.7	90.1*	88.1*	95.2	91.2	88.1*	82.8*
Pregnant women whose abdomen examined	73.3	69.1	98.4	98.8	83.2	85.6	89.9	87.8	71.7	53.4	87.8	81.9
Pregnant women who consumed 100 or more IFA Tablets / Syrup equivalent	58.2	64.9	52.9	49.2	55.5	52.6	44.8	44.3	64.3	60.8	50.8	50.0
Pregnant women who had full antenatal care[5]	57.1	62.8	49.0	45.6	51.1	48.5	39.3	33.7	61.7	56.8	39.3	33.7
Delivery Care (women who had live/still birth during reference period) (%)												
Institutional delivery	92.2	89.4	56.6	54.1	93.2	92.9	56.9	51.0	80.3	68.2	59.8	46.6
Delivery at government health institutions	36.5	44.7	23.2	23.1	52.3	58.2	32.2	32.3	51.9	50.5	31.5	30.5
Delivery at private health institutions	55.6	44.7	34.3	31.3	40.9	34.7	25.5	18.8	28.3	17.7	28.2	15.8
Delivery by Caesarean section at government health institutions	11.2	11.7	4.3	4.4	6.3	7.1	7.5	8.3	12.7	10.4	9.2	6.2
Delivery by Caesarean section at private health institutions	32.8	24.2	10.1	8.3	24.0	16.3	10.9	6.8	19.7	9.9	13.7	6.8
Delivery at home	7.3	9.6	38.2	41.2	6.8	7.1	37.2	43.2	18.8	30.2	39.0	51.7
Delivery at home conducted by skilled health personnel[6] (Out of total Deliveries)	6.7	8.5	13.6	13.8	3.4	3.1	7.4	8.3	3.2	4.7	1.3	1.1
Mothers who received post-natal care within 48 hours of Institutional delivery …	79.4	73.8	75.8	72.9	84.8	87.9	70.3	66.1	86.9	82.4	66.4	56.5
Mothers who received post-natal care within two weeks of Institutional delivery	83.5	77.4	84.1	81.9	89.0	91.2	74.9	70.8	88.8	83.2	72.8	64.6
Delivery attended by skilled health personnel	18.7	19.9	69.8	66.8	19.7	22.2	64.9	59.2	21.2	20.8	61.6	48.9
Discharge of mothers from institution after minimum stay of 48 hours	77.6	79.8	NA	NA	65.2	56.0	NA	NA	74.6	61.1	NA	NA
Out of pocket expenditure per institutional delivery in Public health facility(Rs. in 000's)	3.0	2.9	NA	NA	3.1	3.5	NA	NA	3.8	2.8	NA	NA
Percentage of women who received JSY benefits												
Home delivery	23.1	33.3	16.5	16.0	8.3	14.3	18.0	19.3	6.7	6.8	6.2	4.3
Institutional delivery	19.4	23.8	24.6	28.6	28.7	30.8	21.6	31.6	31.0	38.9	36.2	45.1
Percentage of Women who had												
Any Pregnancy complication[7]	24.6	13.7	52.4	50.5	41.1	34.7	59.0	54.7	36.0	29.7	60.2	54.8
Any Delivery complication[7]	11.7	8.5	64.6	65.7	26.3	31.6	61.3	61.5	15.6	10.4	44.6	45.2
Any Post-delivery complication[7]	7.9	6.4	41.3	40.1	30.7	31.6	40.2	40.1	13.7	9.9	48.6	52.8
Problem of vaginal discharge during last three months	10.5	12.5	11.4	11.9	9.1	10.0	8.3	8.0	8.8	13.5	9.6	13.0
Menstrual related problems during last three months**	19.8	19.0	23.1	23.4	11.0	12.3	21.8	22.1	15.1	20.6	27.2	28.2
Percentage of pregnancy resulted in												
Live Birth	96.9	97.8	93.5	92.7	99.1	98.4	90.3	91.1	98.4	98.2	92.4	94.0
Still Birth	0.8	NA	1.0	1.2	NA	NA	1.6	1.2	0.5	0.7	1.6	1.9
Induced abortion	NA	NA	1.7	1.9	NA	NA	1.0	1.2	0.2	0.4	0.6	NA
Spontaneous abortion	2.4	2.2	3.8	4.2	0.9	1.6	7.1	6.5	0.9	0.7	5.4	4.2

Indicators	East Godavari				West Godavari				Krishna			
	DLHS-4 (2012-13)		DLHS-3 (2007-08)		DLHS-4 (2012-13)		DLHS-3 (2007-08)		DLHS-4 (2012-13)		DLHS-3 (2007-08)	
	TOTAL	RURAL	TOTAL	RURAL	TOTAL	RURAL	TOTAL	RURAL	TOTAL	RURAL	TOTAL	RURAL
Sample Size												
Ever married women age15-49 years	1578	994	869	670	1075	594	814	670	1189	654	1061	745
Population and household profile												
Sex ratio at birth (Male per 100 Female)	98.4	98.8	98.9	99.6	96.9	99.9	101.0	103.2	99.5	97.9	99.4	98.8
Percentage of households												
Improved source of drinking water[1]	98.9	98.3	96.7	95.6	99.2	98.8	98.1	97.7	97.8	96.3	93.4	90.7
Having access to improved toilet facility	69.4	54.9	50.4	39.3	80.3	69.0	56.7	49.9	78.9	65.6	60.3	45.3
Marriage												
Mean age at marriage for girls (marriages that occurred during the reference period)	20.4	20.3	19.1	21.0	19.6	19.5	18.8	18.7	19.3	19.4	19.5	18.6
Mean age at marriage for boys (marriages that occurred during the reference period)	24.4	23.9	23.2	22.4	22.9	21.4	23.8	23.4	23.2	24.1	23.8	23.1
Percentage of currently married women married below age 18 years (marriages that occurred during the reference period)	12.7	13.3	28.6	29.3	14.7	11.0	28.4	25.7	14.0	13.8	20.6	24.7
Characteristics of women (%)												
Currently married women who are illiterate	42.6	46.9	30.3	34.4	36.9	41.0	34.1	35.2	38.5	50.1	31.5	37.8
Currently married women with 10 or more years of schooling	34.2	19.4	15.4	8.1	29.1	23.5	16.8	15.3	35.8	23.1	23.1	14.9
Fertility (%)												
Births to women aged 15-19 years out of total births[3]	7.6	8.4	17.9	19.2	5.6	4.0	16.5	17.4	4.8	5.1	11.0	13.9
Women aged 20-24 years reporting birth order of 2 & above	46.2	44.4	51.6	53.8	55.6	53.1	61.3	66.9	52.8	60.8	47.6	49.6
Women aged 15-49 years who reported birth order of 3 & above	7.6	7.9	11.6	13.2	6.8	8.4	11.0	13.0	6.8	7.2	10.0	9.9
Women with two children wanting no more children	46.8	42.4	22.3	20.1	46.8	42.4	27.2	19.6	25.4	30.6	21.1	16.6
Mean no. of children ever born to women age 40-49 years	2.3	2.3	3.2	3.5	2.3	2.3	3.0	3.1	2.1	2.1	3.0	3.1
Current use of Family Planning Methods (%)												
Any method	71.0	69.4	78.5	77.2	70.9	72.2	78.5	78.8	76.5	78.7	76.4	74.2
Any modern method	70.8	69.0	78.0	76.6	70.8	72.2	78.5	78.9	76.3	78.4	76.4	74.2
Female sterilization	69.3	71.7	74.8	76.1	74.9	77.9	70.8	71.1	73.2	75.4	69.8	72.3
Male sterilization	1.4	0.5	2.8	2.5	1.3	0.3	3.9	2.6	0.4	0.2	0.5	0.3
Pill	0.2	NA	0.7	0.2	NA	NA	3.2	0.1	0.1	0.1	0.6	0.1
IUD	NA	NA	0.3	0.0	0.2	NA	0.8	0.0	0.2	NA	0.8	0.0
Condom	0.1	NA	0.1	0.0	0.5	0.3	0.5	0.3	1.1	0.3	0.6	0.7
Any Traditional method	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Unmet Need for Family Planning (%)												
Total unmet need[4]	19.5	17.4	4.1	3.2	9.3	8.7	6.8	7.1	13.2	11.7	7.2	6.8
Unmet Need for spacing	7.7	6.6	4.7	5.5	11.7	10.6	4.0	4.5	17.9	17.4	3.4	3.6
Unmet Need for limiting	1.6	2.1	2.1	1.6	1.5	1.1	3.2	2.3	1.8	1.7	3.7	3.3
Quality of Family Planning Services (%)												
Non-users ever advised by health personnel to adopt any family planning method	37.7	30.4	15.7	14.7	27.2	20.4	7.9	8.3	16.5	17.6	11.7	12.6
Current users told about side effects of method	6.1	5.9	NA	NA	14.5	6.9	0.0	0.0	11.1	8.7	31.6	33.3
Users who received follow-up services for sterilization and IUD within 48 hours	74.8	82.4	94.1	94.1	76.0	71.5	96.3	96.1	82.9	82.0	91.6	91.8
Post-partum adoption of Family Planning for sterilization	94.5	97.6	94.9	94.7	96.0	93.7	94.4	94.1	87.8	92.0	93.3	94.1
Antenatal Care[5] (Women who had last live/still birth during reference period) (%)												
Pregnant women who received any antenatal check-up	99.2	99.2	97.4	96.9	99.1	99.2	98.9	98.6	100.0	100.0	99.6	99.4
Pregnant women who had antenatal check-up in first trimester	77.3	72.6	53.8	49.6	85.3	78.1	64.8	62.5	92.5	89.8	76.6	75.9
Pregnant women who had three or more ANC visits	93.1	91.6	89.8	91.3	85.0	87.5	95.2	95.9	98.3	99.1	98.3	98.3
Pregnant women who had at least one tetanus toxoid injection	95.7	95.5	94.9	95.2	93.0	92.4	98.1	97.9	96.4	97.3	98.2	98.3
Pregnant women whose Blood Pressure (BP) is taken	91.4	95.1	97.3	96.6	78.6	74.8	98.8	98.5	90.6	86.6	99.6	99.4
Pregnant women who had blood tested (Hb)	84.9	82.9	97.3*	96.7*	78.3	75.8	98.8*	98.5*	84.5	85.8	99.6*	99.4*
Pregnant women whose abdomen examined	64.3	66.1	97.4	96.7	49.6	40.3	98.8	98.5	84.4	81.3	99.5	99.4
Pregnant women who consumed 100 or more IFA Tablets / Syrup equivalent	62.2	65.2	30.3	27.3	45.0	52.1	39.6	38.6	41.7	42.0	42.4	43.1
Pregnant women who had full antenatal care[5]	56.2	57.1	25.6	21.9	39.3	47.4	38.1	36.8	40.1	41.1	59.1	58.6
Delivery Care (women who had live/still birth during reference period) (%)												
Institutional delivery	95.2	94.3	86.6	84.3	97.8	96.7	81.5	82.1	96.9	95.6	84	78.7
Delivery at government health institutions	45.6	43.5	38.5	41.7	45.5	45.0	35.0	37.2	43.5	47.8	37.3	37.7
Delivery at private health institutions	49.6	50.8	48.1	42.5	52.4	51.7	47.0	44.8	53.4	47.8	46.6	41.1
Delivery by Caesarean section at government health institutions	8.6	6.9	5.7	7.0	18.6	19	11.0	11.1	13	14.2	5.9	4.6
Delivery by Caesarean section at private health institutions	26.3	24.8	18.1	12.6	32.3	37.2	21.3	19.3	29.7	30.4	21.6	15.5
Delivery at home	4.5	5.7	11.5	13.4	2.2	3.3	13.8	12.5	3.1	4.4	16.2	21.3
Delivery at home conducted by skilled health personnel[6] (Out of total Deliveries)	1.3	1.6	5.6	6.3	2.2	3.3	4.4	4.8	0.5	0.9	3.9	5.2
Mothers who received post-natal care within 48 hours of Institutional delivery …	70.9	74.6	92.9	92.1	74.8	75.0	89.0	91.0	82.9	76.9	92.4	95.4
Mothers who received post-natal care within two weeks of Institutional delivery	73.5	75.9	94.2	93.7	77.5	78.6	90.1	91.7	87.2	83.3	96.2	98.9
Delivery attended by skilled health personnel	26.3	24.8	18.7	12.6	32.3	37.2	21.3	19.3	29.7	30.4	21.6	15.5
Discharge of mothers from institution after minimum stay of 48 hours	80.6	85.3	NA	NA	85.8	86.2	NA	NA	79.3	83.3	NA	NA
Out of pocket expenditure per institutional delivery in Public health facility(Rs. in 000's)	2.9	2.8	NA	NA	3.0	2.5	NA	NA	3.4	2.8	NA	NA
Percentage of women who received JSY benefits												
Home delivery	5.9	NA	22.2	23.5	NA	NA	7.7	5.6	NA	NA	10.8	13.2
Institutional delivery	23.4	22.4	29.6	33.6	21.7	17.2	27.5	31.1	27.8	30.6	19.6	22.5
Percentage of Women who had												
Any Pregnancy complication[7]	38.4	40.2	67.7	70.9	42.4	39.2	50.3	48.3	47.4	39.3	48.3	46.0
Any Delivery complication[7]	23.1	32.5	59.6	62.2	29.4	19.8	54.9	57.2	25.4	23.0	44.1	40.2
Any Post-delivery complication[7]	19.0	25.2	41.4	44.5	18.2	16.5	23.1	22.2	31.8	23.2	29.2	28.2
Problem of vaginal discharge during last three months	4.5	5.0	11.5	11.7	2.7	1.7	7.3	6.8	12.4	4.6	5.5	5.1
Menstrual related problems during last three months**	11.1	11.3	26.9	27.6	12.9	11.9	23.4	24.7	10.1	4.9	20.9	20.9
Percentage of pregnancy resulted in												
Live Birth	98.6	98.8	97.5	97.1	95.9	97.7	95.2	94.0	100	100	97.1	97.4
Still Birth	0.6	0.6	0.0	0.0	1.2	1.2	0.4	0.5	NA	NA	0.3	0.4
Induced abortion	1.8	NA	NA	0.6	1.8	NA	NA	NA	NA	NA	0.0	0.0
Spontaneous abortion	0.8	0.8	2.0	2.4	1.2	1.2	4.4	5.4	NA	NA	2.6	2.2

Indicators	Guntur				Prakasam				SPSR Nellore			
	DLHS-4 (2012-13)		DLHS-3 (2007-08)		DLHS-4 (2012-13)		DLHS-3 (2007-08)		DLHS-4 (2012-13)		DLHS-3 (2007-08)	
	TOTAL	RURAL	TOTAL	RURAL	TOTAL	RURAL	TOTAL	RURAL	TOTAL	RURAL	TOTAL	RURAL
Sample Size												
Ever married women age15-49 years	1632	1050	1055	759	933	455	833	690	1346	882	844	671
Population and household profile												
Sex ratio at birth (Male per 100 Female)	96.6	98.2	100.3	98.8	99.0	98.0	98.3	99.4	97.1	98.1	100.5	99.3
Percentage of households												
Improved source of drinking water[1]	97.6	98.0	93.2	91.0	99.0	98.0	91.3	90.2	99.1	99.5	88.4	90.0
Having access to improved toilet facility	70.6	58.5	52.2	39.6	70.8	49.4	40.0	31.3	55.9	38.7	38.1	28.4
Marriage												
Mean age at marriage for girls (marriages that occurred during the reference period)	19.3	18.9	18.9	18.4	20.4	19.1	18.4	17.8	20.3	19.3	19.6	19.5
Mean age at marriage for boys (marriages that occurred during the reference period)	23.5	23.3	22.9	22.5	24.4	23.2	22.8	22.7	24.2	23.9	24.9	24.4
Percentage of currently married women married below age 18 years (marriages that occurred during the reference period)	22.1	25.0	30.3	36.5	14.3	23.6	31.7	36.6	14.4	16.3	19.6	21.2
Characteristics of women (%)												
Currently married women who are illiterate	51.6	58.2	43.6	48.8	43.6	58.7	49.7	55.5	34.4	42.2	40.3	44.5
Currently married women with 10 or more years of schooling	23.1	16.4	20.4	13.7	30.9	18.9	17.6	13.4	29.8	22.2	21.3	14.9
Fertility (%)												
Births to women aged 15-19 years out of total births[3]	10.4	13.0	14.6	16.2	6.5	9.8	15.2	17.2	7.2	7.3	13.6	14.8
Women aged 20-24 years reporting birth order of 2 & above	50.2	50.8	51.3	52.9	54.3	55.9	52.1	49.5	58.5	59.8	51.6	51.9
Women aged 15-49 years who reported birth order of 3 & above	12.6	12.1	5.7	3.6	9.6	18.5	15.9	15.8	12.5	14.2	12.1	13.7
Women with two children wanting no more children	24.6	25.7	21.6	16.0	20.3	20.0	25.8	23.3	55.2	52.8	28.1	27.4
Mean no. of children ever born to women age 40-49 years	2.7	2.7	3.0	3.0	2.2	2.6	3.5	3.7	2.4	2.6	2.8	2.9
Current use of Family Planning Methods (%)												
Any method	75.0	76.2	71.8	73.3	72.4	73.1	70.1	69.4	67.5	67.9	66.3	67.0
Any modern method	74.8	76	71.4	73.4	71.6	72.6	70.1	69.7	67.3	67.8	66.1	68.0
Female sterilization	72.3	75.4	69.8	72.3	70.3	72.1	68.3	68.0	67.2	67.9	65	67.2
Male sterilization	0.4	0.2	0.5	0.3	0.5	0.2	0.9	0.5	0.3	NA	1.2	0.2
Pill	0.1	0.1	0.6	0.1	0.5	NA	0.3	0.3	0.1	NA	0.0	0.0
IUD	0.2	0.2	0.1	0.1	NA	NA	0.5	0.5	0.3	0.2	0.4	0.0
Condom	1.1	0.3	0.6	0.7	0.7	0.2	0.4	0.5	NA	NA	0.6	0.5
Any Traditional method	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Unmet Need for Family Planning (%)												
Total unmet need[4]	13.2	11.7	7.2	6.8	19.7	19.1	7.1	6.9	12.6	12.3	7.3	7.7
Unmet Need for spacing	11.7	10.6	4.0	4.5	17.9	17.4	3.4	3.6	7.4	8.0	2.3	2.3
Unmet Need for limiting	1.5	1.1	3.2	2.3	1.8	1.7	3.7	3.3	5.2	4.3	5.0	5.4
Quality of Family Planning Services (%)												
Non-users ever advised by health personnel to adopt any family planning method	16.5	17.6	11.7	12.6	25.9	34.0	17.0	16.7	9.7	9.1	16.5	18.1
Current users told about side effects of method	11.1	8.7	31.6	33.3	7.5	8.8	37.5	42.9	4.3	5.4	NA	NA
Users who received follow-up services for sterilization and IUD within 48 hours	82.9	82.0	91.6	91.8	46.6	48.5	91.0	90.2	60.1	61.7	92.6	93.2
Post-partum adoption of Family Planning for sterilization	95.8	97.0	89.2	91.2	92.8	90.3	75.9	77.7	97.8	98.6	96.3	96.9
Antenatal Care[5] (Women who had last live/still birth during reference period) (%)												
Pregnant women who received any antenatal check-up	98.1	98.8	98.2	97.4	95.0	92.3	95.1	94.3	98.9	99.2	97.3	97.4
Pregnant women who had antenatal check-up in first trimester	85.1	87.0	68.1	64.5	88.7	88.2	62.9	60.4	84.2	83.3	71.8	69.9
Pregnant women who had three or more ANC visits	93.0	92.6	88.0	84.5	86.8	87.7	84.6	81.1	91.6	89.9	71.8	69.9
Pregnant women who had at least one tetanus toxoid injection	95.8	95.5	97.0	96.1	99.0	97.6	95.1	94.3	97.2	98.0	92.5	91.4
Pregnant women whose Blood Pressure (BP) is taken	95.5	96.0	98.0	97.1	98.6	97.6	94.6	93.6	92.6	93.5	97.2	97.3
Pregnant women who had blood tested (Hb)	91.9	91.5	98.0*	97.2*	97.6	96.5	94.8*	93.9*	92.6	92.3	97.2*	97.3*
Pregnant women whose abdomen examined	81.7	80.6	97.9	97.1	97.6	98.8	94.5	93.6	76.3	76.9	94.5	93.6
Pregnant women who consumed 100 or more IFA Tablets / Syrup equivalent	49.7	45.7	43.8	40.0	55.8	50.0	43.5	41.5	45.2	40.5	46.0	43.8
Pregnant women who had full antenatal care[5]	43.8	40.2	41.9	37.4	48.5	43.9	37.0	34.0	39.0	35.1	40.6	37.9
Delivery Care (women who had live/still birth during reference period) (%)												
Institutional delivery	92.1	91.2	77.8	76.8	92.7	91.2	76.1	73.6	91.6	92.4	88.4	87.0
Delivery at government health institutions	29.3	33.3	24.9	27.7	26.4	35.2	24.3	24.5	25.5	27.2	24.9	29.2
Delivery at private health institutions	62.7	57.8	54.4	49.0	66.4	56.0	51.9	49.

Indicators	Srikakulam				Vizianagaram				Visakhapatnam				East Godavari				West Godavari				Krishna			
	DLHS-4 (2012-13)		DLHS-3 (2007-08)		DLHS-4 (2012-13)		DLHS-3 (2007-08)		DLHS-4 (2012-13)		DLHS-3 (2007-08)		DLHS-4 (2012-13)		DLHS-3 (2007-08)		DLHS-4 (2012-13)		DLHS-3 (2007-08)		DLHS-4 (2012-13)		DLHS-3 (2007-08)	
	TOTAL	RURAL	TOTAL	RURAL	TOTAL	RURAL	TOTAL	RURAL	TOTAL	RURAL	TOTAL	RURAL	TOTAL	RURAL	TOTAL	RURAL	TOTAL	RURAL	TOTAL	RURAL	TOTAL	RURAL	TOTAL	RURAL
Child Immunization (%) (Children age 12-23 months)																								
Received full vaccination[8]	80.0	94.7	62.3	58.3	63.4	63.6	55.4	47.5	65.0	59.2	60.0	50.0	62.2	73.6	65.4	61.4	51.6	54.5	72	73	46.7	48.3	82.9	80.0
Received BCG vaccine	95.5	100.0	97.6	98	97.6	95.5	97.4	96.7	95.1	94.0	86.9	81.6	100.0	100.0	99.8	100.0	95.3	93.9	100	100	89.1	86.7	98.6	98.5
Received 3 doses of DPT vaccine	93.2	100.0	67.7	65.3	76.2	81.8	65.5	58.3	85.2	81.6	67.6	60.5	79.3	81.1	72.1	67.4	71.4	84.4	79.1	78.9	51.1	51.7	93.2	92.3
Received 3 doses of polio vaccine	88.9	100.0	73.3	71.4	90.5	87.0	78.2	76.7	81.7	76.0	77.1	76.3	75.6	90.6	81.5	81.4	71.4	66.7	83.5	81.1	69.6	66.7	89.2	87.7
Received measles vaccine	90.9	94.7	90.1	89.8	85.7	82.6	82.2	78.7	91.4	91.8	73.9	68.4	96.4	94.4	90.0	88.4	85.7	87.9	97.8	97.4	91.3	90.0	96.9	95.5
Children (age 9-35 months) received at least one dose of vitamin A supplement in last 6 months	80.5	95.0	NA	NA	80.7	80.0	81.7	79.2	78.1	80.9	65.0	59.4	83.0	81.8	72.8	75.0	88.6	83.3	83.9	84.1	90.4	89.6	87.9	89.3
Child feeding practices (based on last-born child in the reference period) (%)																								
Children age 0-5 months exclusively breastfed[9]	81.3	76.9	NA	NA	75.0	75.0	NA	NA	89.3	88.9	NA	NA	78.8	82.4	NA	NA	80.0	93.3	NA	NA	69.2	62.5	NA	NA
Children age 6-9 months receiving solid/semi-solid food and breast milk	68.8	66.7	50.0	42.9	72.2	78.6	50.0	37.5	78.9	71.4	54.3	50.0	54.8	58.3	35.3	35.3	78.6	75.0	41.2	46.7	50.0	50.0	48.4	50.0
Children age 12-23 months receiving breast feeding along with complementary feeding	72.1	73.7	NA	NA	39.0	36.4	NA	NA	63.8	65.3	NA	NA	58.0	71.7	NA	NA	60.9	50	NA	NA	42.2	44.8	NA	NA
Children age 6-35 months exclusively breastfed for at least 6 months	25.5	24.5	17.0	16.3	22.2	16.1	36.6	38.6	32.1	33.0	28.0	36.4	22.2	21.4	54.1	56.5	10.3	11.1	39.0	42.4	12.5	11.7	32.1	34.1
Children under 3 years breastfed within one hour of birth	41.5	41.9	55.0	53.6	68.1	63.2	59.7	62.7	46.1	47.1	54.2	60.0	50.2	57.5	65.2	68.1	60.3	60.9	53.3	54.2	67.9	63.2	52.5	53.3
Birth Weight (%) (age below 36 months)																								
Percentage of Children weighed at birth	98.3	98.4	NA	NA	99.2	98.5	NA	NA	82.2	73.1	NA	NA	95.5	96.3	NA	NA	92.4	92.0	NA	NA	97.2	97.0	NA	NA
Percentage of Children with low birth weight (out of those who weighted) (below 2.5 kg)	4.3	4.9	NA	NA	5.0	1.5	NA	NA	4.2	2.1	NA	NA	3.0	3.2	NA	NA	9.0	11.1	NA	NA	15.0	12.1	NA	NA
Awareness about Diarrhoea (%)																								
Women know about what to do when a child gets diarrhoea	50.9	64.1	70.7	69.4	79.3	76.3	71.4	69.2	65.0	59.6	68.7	58.8	78.8	76.3	69.4	69.9	75.8	73.4	69.6	68.5	83.7	79.9	77.9	72.6
Awareness about ARI (%)																								
Women aware about danger signs of ARI[10]	39.4	48.5	30.5	31.1	71.2	62.8	29.6	25.9	45.9	41.2	28.6	17.4	67.5	64.8	21.9	20.6	75.4	74.1	32.4	31.3	75.7	71.5	35.2	27.8
Utilization of Government Health Services (%)																								
Antenatal care	45.1	55.9	46.3	45.6	63.0	61.9	56.2	58.5	64.7	68.9	66.8	74.8	50.1	46.5	64.9	74.0	51.1	58.0	44.4	43.7	45.6	50.4	44.0	49.1
Treatment for pregnancy complications	23.5	40.0	35.5	36.7	68.3	72.7	40.8	46.7	56.0	60.0	51.9	68.9	28.6	29.6	42.7	44.2	49.3	35.1	28.6	27.0	39.7	40.0	36.4	40.0
Treatment for post-delivery complications	30.8	40.0	28.3	26.9	59.4	72.2	43.3	46.0	51.7	63.6	46.8	55.2	41.5	41.2	40.5	45.2	50.0	58.8	46.4	57.9	38.1	53.8	33.3	30.8
Nutritional Status of children below 5 years																								
Children below 5 years wasting (weight for height-below 2 SD)	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Children below 5 years wasting (weight for height-below 3 SD)	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Children below 5 years stunting (height for age-below 2 SD)	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Children below 5 years stunting (height for age-below 3 SD)	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Children below 5 years underweight (weight for age-below 2 SD)	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Children below 5 years underweight (weight for age-below 3 SD)	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Anaemia Status by Haemoglobin Level14 (%)																								
Adolescents (15-19 years) having anemia	61.5	67.7	NA	NA	56.4	54.0	NA	NA	52.9	53.0	NA	NA	62.2	62.5	NA	NA	61.1	53.3	NA	NA	67.5	71.8	NA	NA
Adolescents (15-19 years) having severe anemia	4.6	6.9	NA	NA	5.5	6.0	NA	NA	4.9	7.4	NA	NA	7.8	9.0	NA	NA	8.9	10.9	NA	NA	11.6	13.5	NA	NA
Pregnant women (15-49 aged) having anemia	78.0	86.5	NA	NA	71.1	71.4	NA	NA	61.8	64.7	NA	NA	63.9	65.1	NA	NA	60.9	59.0	NA	NA	82.4	77.8	NA	NA
Pregnant women (15-49 aged) having severe anemia	5.1	5.4	NA	NA	17.8	23.8	NA	NA	8.0	3.9	NA	NA	9.1	9.2	NA	NA	12.0	11.5	NA	NA	15.6	19.2	NA	NA
Women (15-49 aged) having anemia	69.4	75.6	NA	NA	60.4	61.8	NA	NA	61.2	62.4	NA	NA	72.8	71.6	NA	NA	66.7	57.6	NA	NA	72.8	74.6	NA	NA
Women (15-49 aged) having severe anemia	7.5	9.3	NA	NA	5.0	5.6	NA	NA	6.6	7.9	NA	NA	8.2	7.7	NA	NA	8.8	6.5	NA	NA	12.3	16.7	NA	NA
Health programmes at village level																								
Percentage of villages having ASHA	96.0		59.0		100.0		68.2		93.6		63.3		88.6		71.0		88.0		82.5		93.9		79.4	
Percentage of Villages having Village Health Nutrition and Sanitation Committee (VHNSC)	84.0		36.3		60.0		53.6		71.0		53.3		79.6		60.5		92.0		95.0		91.2		44.1	
Accessibility of health facility (%)																								
Villages with Sub-Health Centre within 3 km	92.0		79.5		88.0		67.5		87.1		46.6		95.5		80.5		88.0		87.1		88.2		78.1	
Villages with PHC within 10 km	84.0		77.2		84.0		65.0		80.7		66.6		86.4		63.1		84.0		82.5		58.8		79.4	
Availability of Health Infrastructure, Staff and Services at (%)																								
Sub-Health Centre																								
Sub-Health Centre located in government building	8.3		11.5		24.0		12.8		22.6		29		22.7		21.4		16.0		21.4		8.8		21.1	
Sub-Health Centre with ANM	100.0		90.0		84.0		89.7		100.0		80.7		97.7		83.3		88.0		88.1		91.2		81.6	
Sub-Health Centre with male health worker	58.3		40.0		44.0		66.66		9.7		29.0		50.0		69.0		44.0		50.0		8.8		44.7	
Sub-Health Centre with ANM residing in Sub-Health Centre quarter where facility is available	50.0		100.0		100.0		66.7		100.0		50.0		100.0		62.5		NA		100.0		100.0		100.0	
Sub-Health Centre with additional ANM	79.2		42.5		71.4		69.2		54.8		29.0		83.7		0.0		63.6		11.9		64.5		21.1	
Primary Health Centre (PHC)																								
PHCs functioning on 24 X 7 hours basis	42.9		31.0		40.0		52.0		63.0		50.0		21.9		42.8		35.0		72.7		39.3		45.8	
PHCs having Lady Medical Officer	63.2		31.0		47.1		40.0		34.6		37.5		46.2		32.1		31.6		27.2		60.7		25	
PHCs having residential quarter for Medical Officer	9.5		20.6		15.0		12.0		14.8		12.5		3.1		17.8		20.0		24.2		10.7		4.1	
PHCs having new born care services on 24 X 7 hours basis	100.0		NA		100.0		NA		100.0		NA		100		NA		100.0		NA		92.9		NA	
PHCs having referral services for pregnancies/delivery on 24 X 7 hours basis	66.7		NA		87.5		NA		64.7		NA		71.4		NA		71.4		NA		54.6		NA	
PHCs conducted at least 10 deliveries during last one month on 24 X 7 hours basis	72.7		41.4		63.6		72.0		75.0		45.8		38.5		64.3		25.0		54.6		21.4		54.2	

Indicators	Guntur				Prakasam				SPSR Nellore				Chittoor				YSR Kadapa				Ananthapur				Kurnool				Andhra Pradesh					
	DLHS-4 (2012-13)		DLHS-3 (2007-08)		DLHS-4 (2012-13)		DLHS-3 (2007-08)		DLHS-4 (2012-13)		DLHS-3 (2007-08)		DLHS-4 (2012-13)		DLHS-3 (2007-08)		DLHS-4 (2012-13)		DLHS-3 (2007-08)		DLHS-4 (2012-13)		DLHS-3 (2007-08)		DLHS-4 (2012-13)		DLHS-3 (2007-08)		DLHS-4 (2012-13)		DLHS-3 (2007-08)			
	TOTAL	RURAL	TOTAL	RURAL	TOTAL	RURAL	TOTAL	RURAL	TOTAL	RURAL	TOTAL	RURAL	TOTAL	RURAL	TOTAL	RURAL	TOTAL	RURAL	TOTAL	RURAL	TOTAL	RURAL	TOTAL	RURAL	TOTAL	RURAL	TOTAL	RURAL	TOTAL	RURAL	URBAN	TOTAL	RURAL	URBAN
Child Immunization (%) (Children age 12-23 months)																																		
Received full vaccination[8]	62.8	66.1	59.4	54.7	57.1	55	58.5	61.1	54.4	56.4	55.6	56.8	55.3	61.5	62.0	63.6	62.0	63.7	64.1	62.2	66.2	66.7	71.2	68.6	59.4	61.0	54.7	46.3	55.4	57.9	51.0	67.1	65	73.2
Received BCG vaccine	97.4	96.4	97.3	96.2	91.8	84.2	100.0	100.0	96.2	96.4	96.5	95.5	94.8	92.5	99.4	100.0	97.2	98.5	98.2	100.0	97.4	96.6	97.9	97.1	94.3	93.4	97.7	97.6	2.1	2.3	1.8	1.0	1.1	0.8
Received 3 doses of DPT vaccine	80.8	80.4	77.9	77.4	72.0	80.0	62.0	64.8	67.9	70.4	78.3	77.3	66.2	69.8	70.7	69.7	81.7	81.5	85.0	80.0	81.8	79.7	92.8	91.2	72.6	75.3	68.3	64.2	94.1	93.7	94.8	97.5	96.9	99.0
Received 3 doses of polio vaccine	82.1	83.9	73.0	71.2	92.0	85.0	86.7	87.0	93.7	94.5	69.7	70.5	81.3	80.8	80.0	79.4	76.9	78.5	78.8	77.3	81.8	81.4	85.4	88.2	79.0	80.3	80.2	77.8	71.1	73.4	66.7	79.0	77.0	85.2
Received measles vaccine	80.8	78.6	92.8	92.5	79.6	70.0	94.1	92.6	84.8	83.6	87.6	88.6	86.8	86.5	88.5	90.0	83.3	85.9	82.0	82.2	84.4	84.7	94.7	91.4	86.7	86.8	84.9	82.7	79.3	80.0	77.8	82.1	81.4	84.1
Children (age 9-35 months) received at least one dose of vitamin A supplement in last 6 months	78.3	73.8	73.1	72.2	84.5	75.6	80.5	79.4	85.6	84.3	75.9	74.2	86.8	88.3	86.3	90.1	77.1	86.2	76.6	76.0	85.8	87.4	79.5	78.3	77.1	75.5	81.7	79.3	81.6	82.0	80.9	78.8	77.9	81.7
Child feeding practices (based on last-born child in the reference period) (%)																																		
Children age 0-5 months exclusively breastfed[9]	60.0	65.6	NA	NA	72.4	66.7	NA	NA	74.5	68.8	NA	NA	65.8	59.1	NA	NA	65.5	63.4	NA	NA	81.1	79.3	NA	NA	79.2	84.7	NA	NA	70.1	70.2	69.8	43.9	46.2	36.4
Children age 6-9 months receiving solid/semi-solid food and breast milk	73.0	78.3	65.2	66.7	62.5	60.0	90.5	93.8	66.7	66.7	45.5	43.8	42.9	45.0	60.0	56.3	64.6	67.7	60.0	52.6	54.4	54.8	34.4	30.8	65.5	68.3	36.0	25.0	62.8	64.4	59.3	55	52.3	63.1
Children age 12-23 months receiving breast feeding along with complementary feeding	70.1	65.5	NA	NA	44.9	45.0	NA	NA	51.3	46.3	NA	NA	42.9	37.0	NA	NA	54.2	55.4	NA	NA	41.8	47.5	NA	NA	51.9	44.6	NA	NA	53.1	52.4	54.3	NA	NA	NA
Children age 6-35 months exclusively breastfed for at least 6 months	30.7	28.0	29.3	35.1	30.0	30.4	39.3	43.5	21.7	21.4	35.5	37.1	31.0	25.0	37.9	41.9	24.8	19.1	37.7	45.8	20.4	19.5	34.3	33.3	22.9	20.2	48.5	50.7	21.7	20.6	23.7	32.3	34.7	25.3
Children under 3 years breastfed within one hour of birth	49.2	45.6	51.6	51.5	59.0	55.2	40.6	40.4	54.0	56.7	53.5	53.7	68.7	66.2	62.8	60.7	46.1	41.8	45.3	42.3	58.6	59.9	54.0	55.3	52.1	55.7	53.0	52.0	54.3	54.4	54.3	47.8	49.6	42.4
Birth Weight (%) (age below 36 months)																																		
Percentage of Children weighed at birth	97.5	97.5	NA	NA	95.7	94.6	NA	NA	93.2	93.0	NA	NA	91.4	89.4	NA	NA	96.1	96.0	NA	NA	87.8	83.5	NA	NA	81.4	78.5	NA	NA	91.2	89.6	94.5	NA	NA	NA
Percentage of Children with low birth weight (out of those who weighted) (below 2.5 kg)	7.8	9.1	NA	NA	2.3	NA	NA	NA	5.0	5.0	NA	NA	8.3	9.5	NA	NA	6.6	4.1	NA	NA	4.0	4.6	NA	NA	4.1	4.3	NA	NA	6.9	6.4	7.8	NA	NA	NA
Awareness about Diarrhoea (%)																																		
Women know about what to do when a child gets diarrhoea	58.1	53.7	80.1	79.2	70.4	64.4	63.4	63.5	55.4	52.1	77.8	76.3	79.7	78.6	57.1	49.6	62.1	60.5	73.5	70.4	69.1	64.8	77.2	77.1	46.1	42.2	55.8	62.7	68.8	66.3	73.4	70.4	67.7	78.4
Awareness about ARI (%)																																		
Women aware about danger signs of ARI[10]	47.6	41.5	33.6	33.2	67.9	68.2	34.0	35.3	49.2	44.6	32.2	31.0	70.8	64.3	28.0	24.2	47.3	43.6	31.0	30.1	56.8	56.8	28.4	24.2	32.0	34.2	26.8	24.2	59.2	56.5	64.2	36.7	32.7	48.8
Utilization of Government Health Services (%)																																		
Antenatal care	37.5	38.5	41.8	45.0	26.3	35.3	40.6	40.7	28.4	28.3	23.1	26.2	64.1	75.9	77.9	72.3	22.0	23.9	36.9	39.5	55.2	59.9	51.3	52.1	45.0	49.7	50.7	53.8	43.1	48.0	33.7	43.9	46.2	37.0
Treatment for pregnancy complications	27.9	34.1	34.1	39.7	37.5	47.1	26.6	27.1	27.2	27.4	32.3	33.3	50.0	81.0	40.4	48.6	12.7	13.6	32.3	33.3	45.9	51.7	23.3	19.3	25.3	23.1	42.4	53.3	35.8	39.7	28.8	31.4	33.0	27.5
Treatment for post-delivery complications	50.0	63.6	16.2	22.7	61.1	57.1	20.8	17.6	26.3	20.7	21.1	33.3	39.3	54.5	70.8	75.0	23.3	17.6	21.1	33.3	31.3	33.3	39.1	47.1	30.4	25.0	25.0	20.8	37.9	41.0	32.4	31.8	33.6	25.7
Nutritional Status of children below 5 years																																		
Children below 5 years wasting (weight for height-below 2 SD)	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	25.9	26	25.7	NA	NA	NA
Children below 5 years wasting (weight for height-below 3 SD)	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	15.3	15.8	14.5	NA	NA	NA
Children below 5 years stunting (height for age-below 2 SD)	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	26.2	26	26.5	NA	NA	NA
Children below 5 years stunting (height for age-below 3 SD)	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	14.6	15.4	13.4	NA	NA	NA
Children below 5 years underweight (weight for age-below 2 SD)	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	28.1	30.1	25.2	NA	NA	NA
Children below 5 years underweight (weight for age-below 3 SD)	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	11.3	12.5	9.4	NA	NA	NA
Anaemia Status by Haemoglobin Level14 (%)																																		
Adolescents (15-19 years) having anemia	63.7	70.8	NA	NA	52.3	50.8	NA	NA	64.6	66.9	NA	NA	56.7	66.1	NA	NA	50.1	52.2	NA	NA	59.5	68.1	NA	NA	63.4	67.3	NA	NA	56.7	57.9	54.2	NA	NA	NA
Adolescents (15-19 years) having severe anemia	8.0	10.8	NA	NA	10.1	10.2	NA	NA	6.8	4.7	NA	NA	6.8	9.6	NA	NA	7.7	6.9	NA	NA	11.7	14.1	NA	NA	10.0	7.7	NA	NA	7.4	7.7	6.6	NA	NA	NA
Pregnant women (15-49 aged) having anemia	79.5	83.0	NA	NA	57.7	46.2	NA	NA	84.6	87.7	NA	NA	70.6	71.7	NA	NA	59.8	59.7	NA	NA	74.7	82.4	NA	NA	73.6	75.0	NA	NA	67.4	67.5	67.0	NA	NA	NA
Pregnant women (15-49 aged) having severe anemia	11.7	14.1	NA	NA	NA	NA	NA	NA	10.8	8.1	NA	NA	10.9	15.0	NA	NA	6.6	6.5	NA	NA	18.3	19.4	NA	NA	9.2	10.6	NA	NA	9.2	9.4	8.6	NA	NA	NA
Women (15-49 aged) having anemia	73.7	77.2	NA	NA	64.7	64.2	NA	NA	72.4	71.2	NA	NA	63.4	66.5	NA	NA	65.4	65.5	NA	NA	63.2	67.9	NA	NA	69.5	71.1	NA	NA	64.0	64.2	63.5	NA	NA	NA
Women (15-49 aged) having severe anemia	9.6	11.0	NA	NA	9.7	10.3	NA	NA	9.6	6.9	NA	NA	8.9	10.5	NA	NA	9.1	9.9	NA	NA	12.4	14.8	NA	NA	12.1	11.2	NA	NA	8.1	8.4	7.6	NA	NA	NA
Health programmes at village level																																		
Percentage of villages having ASHA	90.9		63.8		84.0		92.8		90.9		35.9		86.4		84.2		97.7		71.8		95.5		70.2		88.6		86.8		92.6			63.2		
Percentage of Villages having Village Health Nutrition and Sanitation Committee (VHNSC)	50.0		88.3		72.0		61.9		77.3		66.7		72.7		86.8		93.2		74.4		77.3		50.7		54.6		76.3		75.5			67.4		
Accessibility of health facility (%)																																		
Villages with Sub-Health Centre within 3 km	95.5		82.8		52.0		18.5		79.6		0.0		70.5		65.7		77.3		73.0		84.1		67.6		79.6		69.4		84.2			62.1		
Villages with PHC within 10 km	68.2		68.5		84.0		61.0		65.9		88.2		70.5		68.4		65.9		79.5		79.6		48.6		68.2		50.0		70.7			67.9		
Availability of Health Infrastructure, Staff and Services at (%)																																		
Sub-Health Centre																																		
Sub-Health Centre located in government building	34.1		12.2		12.5		19.5		13.6		21.6		46.5		33.3		25.6		35.4		27.3		48.7		18.6		4.8		26.3			21.9		
Sub-Health Centre with ANM	100.0		89.8		87.5		78.1		79.6		86.5		76.7		78.6		97.7		64.6		81.8		94.9		93.0		69.1		92.8			84.4		
Sub-Health Centre with male health worker	13.6		36.7		33.3		36.7		18.2		59.5		34.9		21.4		69.8		41.6		68.2		64.1		41.9		28.5		33.6			40.6		
Sub-Health Centre with ANM residing in Sub-Health Centre quarter where facility is available	66.7		57.1		--		75.0		100.0		0.0		33.3		75.0		50.0		63.6		20.0		33.3		NA		100.0		58.3			63.3		
Sub-Health Centre with additional ANM	75.0		14.3		66.7		73.2		65.7		2.7		84.9		7.1		90.5		75.0		80.6		76.2		50.0		76.2		72.8			28.0		
Primary Health Centre (PHC)																																		
PHCs functioning on 24 X 7 hours basis	41.4		39.3		33.3		46.2		38.2		40.0		42.4		42.4		48.3		57.1		69.7		75.0		37.8		63.6		47.5			51.1		
PHCs having Lady Medical Officer	46.4		48.4		53.3		38.4		34.4		48.0		31.0		45.4		40.0		28.6		35.5		35.7		32.3		63.6		42.3			40.6		
PHCs having residential quarter for Medical Officer	13.8		24.2		0.0		30.7		8.8		4.0		18.2		30.3		20.7		40.0		18.2		46.4		8.1		27.2		15.7			25.7		
PHCs having new born care services on 24 X 7 hours basis	84.6				90.0		NA		87.5		NA		94.1		NA		81.3		NA		95.8		NA		86.7		NA		92.0			95.8		
PHCs having referral services for pregnancies/delivery on 24 X 7 hours basis	66.7				100.0		NA		92.3		NA		92.9		NA		64.3		NA		82.6		NA		85.7		NA		75.1			72.4		
PHCs conducted at least 10 deliveries during last one month on 24 X 7 hours basis	21.4		42.4		20.0		42.3		18.8		28.0		55.6		45.5		41.2		42.9		63.0		67.9		62.5		57.6		43.8			54.8		